ELSEVIER

Environmental Pollution 135 (2005) I–IV

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 135

Number 1 2005

Research papers

1 **Reduction of silver solubility by humic acid and thiol ligands during acanthite (β-Ag_2S) dissolution**
A.R. Jacobson, C.E. Martínez, M. Spagnuolo, M.B. McBride, P. Baveye

The presence of humic acids and thiol ligands reduced Ag dissolution from acanthite particles with mixed-oxidation surface layers.

11 **Bioaccumulation of PCBs in the sea urchin Paracentrotus lividus: seawater and food exposures to a ^{14}C-radiolabelled congener (PCB#153)**
B. Danis, O. Cotret, J.L. Teyssié, P. Bustamante, S.W. Fowler, M. Warnau

The sea urchin Paracentrotus lividus is a valuable indicator for PCB contamination.

17 **Rapid regional recovery from sulfate and nitrate pollution in streams of the western Czech Republic – comparison to other recovering areas**
V. Majer, P. Krám, J.B. Shanley

Acidified streams experienced unprecedented recovery from sulfate and nitrate pollution but pH and ANC decreased in alkaline streams.

29 **Effects of increased deposition of atmospheric nitrogen on an upland moor: leaching of N species and soil solution chemistry**
M.G. Pilkington, S.J.M. Caporn, J.A. Carroll, N. Cresswell, J.A. Lee, T.W. Ashenden, S.A. Brittain, B. Reynolds, B.A. Emmett

Nitrate leaching from an upland moor podsol was significantly increased only in response to rates of N deposition in excess of 96 kg N ha^{-1} yr^{1}.

41 **Risk-based approach to appraise valve closure in the clam *Corbicula fluminea* in response to waterborne metals**
C.-M. Liao, L.-J. Jou, B.-C. Chen

A model was developed to link valve closure in clams to concentrations of metals in water.

53 **Distribution of rare earth elements in an alluvial aquifer affected by acid mine drainage: the Guadiamar aquifer (SW Spain)**
M. Olías, J.C. Cerón, I. Fernández, J. De la Rosa

Pollution of the aquifer with rare earth elements is documented at a site of a major spill from a mining operation.

65 **Interaction of limestone grains and acidic solutions from the oxidation of pyrite tailings**
M. Simón, F. Martín, I. García, P. Bouza, C. Dorronsoro, J. Aguilar

Basaluminite, schwertmannite and jarosite armored the limestone grains, and almost all trace elements co-precipitated, but the precipitation of As and Pb tended to decrease as the pH rose.

73 **Metallothionein induction, antioxidative responses, glycogen and growth changes in *Tubifex tubifex* (Oligochaete) exposed to the fungicide, fenhexamid**
Y.Y. Mosleh, S. Paris-Palacios, M. Couderchet, S. Biagianti-Risbourg, G. Vernet

Exposure to the fungicide fenhexamid increased metallothionein levels in Tubifex tubifex.

83 **An inventory of nitric oxide emissions from soils in China**
Q. Wang, Z. Han, Y. Higano

Land use and nitrogen fertilizer use affect NO emissions from soil.

91 **Sources of variation in concentrations of nickel and copper in mountain birch foliage near a nickel-copper smelter at Monchegorsk, north-western Russia: results of long-term monitoring**
M.V. Kozlov

Foliar concentrations of nickel and copper did not reflect emission decline during 1991–2003; annual variation was explained by weather conditions.

101 **Isoprene emission from tropical tree species**
P.K. Padhy, C.K. Varshney

Isoprene flux (diurnal and seasonal) from some tropical tree species was estimated and a regional comparison was made.

S0269–7491(05)00095-3

111 Epiphytic lichen diversity on dead and dying conifers under different levels of atmospheric pollution
M. Hauck

Epiphytic lichen diversity differs increasingly between healthy and dead or dying conifers with increasing atmospheric pollutant load.

121 Survival of brown trout during spring flood in DOC-rich streams in northern Sweden: the effect of present acid deposition and modelled pre-industrial water quality
H. Laudon, A.B.S. Poléo, L.A. Vøllestad, K. Bishop

High levels of complexed aluminum, at pH levels below 5.0, predisposes brown trout to sulfur-caused damage in the spring.

131 Seasonal exposures to triazine and other pesticides in surface waters in the western Highveld corn-production region in South Africa
L.H. Du Preez, P.J. Jansen van Rensburg, A.M. Jooste, J.A. Carr, J.P. Giesy, T.S. Gross, R.J. Kendall, E.E. Smith, G. Van Der Kraak, K.R. Solomon

Concentrations of atrazine in an agricultural region vary greatly and are influenced by rainfall amounts.

143 Atmospheric versus biological sources of polycyclic aromatic hydrocarbons (PAHs) in a tropical rain forest environment
M. Krauss, W. Wilcke, C. Martius, A.G. Bandeira, M.V.B. Garcia, W. Amelung

Evidence of non-pyrolytic, biogenic production of PAHs is provided.

155 Evaluation of quick tests for phosphorus determination in dairy manures
A. Lugo-Ospina, T.H. Dao, J.A. Van Kessel, J.B. Reeves

The dissolved phosphorous quick test can provide real-time information on soluble manure nutrient control.

163 Trace elements in loggerhead turtles (*Caretta caretta*) from the eastern Mediterranean Sea: overview and evaluation
M.M. Storelli, A. Storelli, R. D'Addabbo, C. Marano, R. Bruno, G.O. Marcotrigiano

Cadmium concentrations in samples from turtles may be high enough to affect their health.

171 Assessment of municipal waste compost as a daily cover material for odour control at landfill sites
C. Hurst, P. Longhurst, S. Pollard, R. Smith, B. Jefferson, J. Gronow

Practical measures to improve landfill odour control are investigated.

179 Mercury assessment and evaluation of its impact on fish in the Cecina river basin (Tuscany, Italy)
R. Scerbo, T. Ristori, B. Stefanini, S. De Ranieri, C. Barghigiani

Mercury entering water from a chlor-alkali plant near Tuscany has led to contamination of river food webs.

Number 2 2005

Research papers

187 Decomposition in soil microcosms of leaves of the metallophyte *Arabidopsis halleri*: effect of leaf-associated heavy metals on biodegradation
U. Boucher, M. Balabane, I. Lamy, P. Cambier

Decomposition of slowly decomposable fractions of metal-rich plant residues is slowed by the metals.

195 Sub-cellular partitioning of Cd, Cu and Zn in tissues of indigenous unionid bivalves living along a metal exposure gradient and links to metal-induced effects
E. Bonneris, O. Perceval, S. Masson, L. Hare, P.G.C. Campbell

In natural populations of bivalves, metals were differentially bound in the gills and digestive gland.

209 Metal stress consequences on frost hardiness of plants at northern high latitudes: a review and hypothesis
K. Taulavuori, M.N.V. Prasad, E. Taulavuori, K. Laine

Metal stress may reduce plant frost hardiness.

221 Dynamical coupling of PBPK/PD and AUC-based toxicity models for arsenic in tilapia *Oreochromis mossambicus* from blackfoot disease area in Taiwan
C.-M. Liao, H.-M. Liang, B.-C. Chen, S. Singh, J.-W. Tsai, Y.-H. Chou, W.-T. Lin

Integrated toxicity models can identify dynamic responses of fish to arsenic.

235 Octanol-solubility of dissolved and particulate trace metals in contaminated rivers: implications for metal reactivity and availability
A. Turner, E. Mawji

New approaches are presented for fractionating trace metals in natural waters.

245 Dietary mercury exposure and bioaccumulation in amphibian larvae inhabiting Carolina bay wetlands
J.M. Unrine, C.H. Jagoe, A.C. Brinton, H.A. Brant, N.T. Garvin

Tadpoles accumulate significant concentrations of mercury in isolated wetlands.

255 Diversity of arbuscular mycorrhizal fungi in grassland spontaneously developed on area polluted by a fertilizer plant
C. Renker, V. Blanke, F. Buscot

Almost all plants species were mycorrhizal.

267 In vitro toxicity of selected pesticides on RTG-2 and RTL-W1 fish cell lines
M.M. Babín, J.V. Tarazona

In vitro cell cultures can provide sensitive indicators for pesticide effects on biota.

275 Early-phase immunodetection of metallothionein and heat shock proteins in extruded earthworm coelomocytes after dermal exposure to metal ions
J. Homa, E. Olchawa, S.R. Stürzenbaum, A. John Morgan, B. Plytycz

Metals upregulate stress response proteins in earthworm coelomocytes.

281 Oxygen levels *versus* chemical pollutants: do they have similar influence on macrofaunal assemblages? A case study in a harbour with two opposing entrances
J.M. Guerra-García, J.C. García-Gómez

A high biomass of benthic animals was found in a metal-polluted estuary where high oxygen levels occurred.

293 The Solling roof revisited – slow recovery from acidification observed and modeled despite a decade of "clean-rain" treatment
L. Martinson, N. Lamersdorf, P. Warfvinge

Model predictions show slow recovery by 2050.

303 The effect of hydrological regime on the metal bioavailability for the wetland plant species *Salix cinerea*
B. Vandecasteele, P. Quataert, F.M.G. Tack

A hydrological regime aiming at wetland creation is a potential management option that favors reducing Cd plant availability in polluted freshwater wetlands.

313 Manganese toxicity thresholds for restoration grass species
M.W. Paschke, A. Valdecantos, E.F. Redente

Mn phytotoxicity thresholds for restoration grasses should be useful for risk assessments of metal-contaminated lands.

323 Decomposition of leaves of the metallophyte *Arabidopsis halleri* in soil microcosms: fate of Zn and Cd from plant residues
U. Boucher, I. Lamy, P. Cambier, M. Balabane

Changes in metallic speciation occurred in two stages.

333 Effects of arsenic on concentration and distribution of nutrients in the fronds of the arsenic hyperaccumulator *Pteris vittata* L.
C. Tu, L.Q. Ma

Arsenic hyperaccumulator P. vittata maintained adequate levels of essential nutrients in the fronds under arsenic stress.

Letter to the editor

341 Comment on "Characterization of a reference site for quantifying uncertainties related to soil sampling" by S. Barbizzi et al. (2004)
P. Baveye

Number 3 2005

SPECIAL ISSUE SECTION: The National Atmospheric Deposition Program (25th Anniversary) and Ammonia Workshop

343 The National Atmospheric Deposition Program: 25 years of monitoring in support of science and policy: An ammonia workshop: the state of science and future needs
J.A. Lynch, M. Kerchner

347 Spatial and temporal trends of precipitation chemistry in the United States, 1985–2002
C.M.B. Lehmann, V.C. Bowersox, S.M. Larson

Data from the National Atmospheric Deposition Program National Trends Network indicate that significant changes in precipitation chemistry have occurred from 1985 to 2002.

363 Trends in atmospheric ammonium concentrations in relation to atmospheric sulfate and local agriculture
V.R. Kelly, G.M. Lovett, K.C. Weathers, G.E. Likens

Relationships between NH_4^+ and SO_4^{2-} suggest that aerosol NH_4^+ may be more strongly linked to SO_2 emissions than to local agricultural NH_3 emissions.

371 Nonlinear regression and ARIMA models for precipitation chemistry in East Central Florida from 1978 to 1997
D.M. Nickerson, B.C. Madsen

Autoregressive integrated moving average models of precipitation data are an improvement over nonlinear models for the prediction of precipitation chemistry composition.

381 Overview and assessment of techniques to measure ammonia emissions from animal houses: the case of the Netherlands
J. Mosquera, G.J. Monteny, J.W. Erisman

There are no methods to measure ammonia effectively from outdoor stockyards.

389 Measuring ammonia emissions from land applied manure: an intercomparison of commonly used samplers and techniques
T.H. Misselbrook, F.A. Nicholson, B.J. Chambers, R.A. Johnson

Three commonly used ammonia emission measurement techniques were compared for different manure types and gave CVs between 23 and 74%.

399 Managing ammonia emissions from livestock production in Europe
J. Webb, H. Menzi, B.F. Pain, T.H. Misselbrook, U. Dämmgen, H. Hendriks, H. Döhler

Reducing NH_3 emissions following spreading of manures to land ranks highly because of the large abatement potential and relatively small cost.

407 Spatial and temporal variability of the overall error of National Atmospheric Deposition Program measurements determined by the USGS collocated-sampler program, water years 1989–2001
G.A. Wetherbee, N.E. Latysh, J.D. Gordon

Spatial and temporal characteristics and confidence limits associated with the overall absolute error in NADP/NTN measurements are identified and quantified.

419 Deposition monitoring networks: what monitoring is required to give reasonable estimates of ammonia/ammonium?
J.W. Erisman, A. Hensen, J. Mosquera, M. Sutton, D. Fowler

Current measurement methods allow monitoring of the deposition of ammonia and ammonium in a network consisting of core sites and low cost sites.

433 Atmospheric nitrogen inputs to the Delaware Inland Bays: the role of ammonia
J.R. Scudlark, J.A. Jennings, M.J. Roadman, K.B. Savidge, W.J. Ullman

Local emissions from poultry production appear to significantly contribute to wet and dry atmospheric NH_x loading to the Delaware Inland Bays.

445 Improved daily precipitation nitrate and ammonium concentration models for the Chesapeake Bay Watershed
J.W. Grimm, J.A. Lynch

A linear least-squares regression approach was used to develop daily precipitation nitrate and ammonium concentration models for the Chesapeake Bay Watershed.

Research Papers

457 Chemometrics methods for the investigation of methylmercury and total mercury contamination in mollusks samples collected from coastal sites along the Chinese Bohai Sea
W. Yawei, L. Lina, S. Jianbo, J. Guibin

Rapana venosa might be used as a potential biomonitor for Hg pollution in the Bohai Sea, China.

469 Effects of increased deposition of atmospheric nitrogen on an upland *Calluna* moor: N and P transformations
M.G. Pilkington, S.J.M. Caporn, J.A. Carroll, N. Cresswell, J.A. Lee, B.A. Emmett, D. Johnson

There are low rates of nitrogen loss from a moor ecosystem under both ambient and high atmospheric loading of nitrogen.

481 Risk to breeding success of waterbirds by contaminants in Hong Kong: evidence from trace elements in eggs
J.C.W. Lam, S. Tanabe, M.H.W. Lam, P.K.S. Lam

Analysis of trace element concentrations in eggs indicated that mercury and selenium might pose some risk to waterbirds in Hong Kong.

491 Equilibrium partitioning of ^{14}C-benzo(a)pyrene and ^{14}C-benazolin between fractionated phases from an arable topsoil
J.-M. Séquaris, A. Lavorenti, P. Burauel

The partition coefficient (K_d) of chemicals was calculated according to two-phase or three-phase partitioning models.

501 The use of autecological and environmental parameters for establishing the status of lichen vegetation in a baseline study for a long-term monitoring survey
S. Gombert, J. Asta, M.R.D. Seaward

Ecological clustering which establishes the status of lichen vegetation can be reliably used as a baseline study to monitor environmental changes.

515 Soil remediation: humic acids as natural surfactants in the washings of highly contaminated soils
P. Conte, A. Agretto, R. Spaccini, A. Piccolo

Solutions of natural humic acids appear to be a better choice for washing highly polluted soils.